

Biochemical profiling of antifungal activity of betel leaf (*Piper betle L.*) extract and its significance in traditional medicine

Sarika Pawar^{1#}, Vidya Kalyankar², Bela Dhamangaonkar¹, Sharada Dagade³,
Shobha Waghmode^{2*} and Abhishek Cukkemane¹

¹Bijasu Agri Research Laboratory LLP, Sr. No. 37, Kondhwa Industrial Estate, Khadi Machine Chowk, Kondhwa, Pune-411048, India

²Department of Chemistry, M. E. S Abasaheb Garware College, Pune - 411 007, Maharashtra, India

³Department of Chemistry, Y. M. College, Bharathi Vidyapeeth, Pune-411038, Maharashtra, India

#Present Address: Microbial Diversity Research Centre, Dr. D. Y. Patil Biotechnology and Bioinformatics Institute, Dr. D. Y. Patil Vidyapeeth, Tathawade, Pune-411033, India

Received: 13 February, 2017; Accepted: 10 March, 2017; Published: 20 March, 2017

*Corresponding author: ShobhaWaghmode, Department of Chemistry, M. E. S Abasaheb Garware College, Pune - 411 007, Maharashtra, India; Tel. No: +91-9422-31-6985; E-mail: shobhawaghmode@yahoo.co.in

Abstract

Piper betle (Linn) commonly called as betel leaf is a widely cultivated plant in the Indian subcontinent. The traditional Indian ayurvedic document describes several of its medicinal properties including as an effective antifungal agent. The present study was conducted to evaluate the secondary metabolite that contributes to its antifungal activity. *In vitro* studies were performed on molds and yeasts on antifungal using fractions obtained from ethyl acetate, hexane and ethanol-methanol extracts by well-diffusion technique. Ethyl acetate extracts showed highest anti-fungal activity. Using biophysical techniques such as Nuclear Magnetic Resonance and Fourier transform infrared spectroscopy techniques; we identified the molecule as derivative of the phenyl propanoid family akin eugenol. The molecule can be readily purified using a 2 step solvent extraction procedure along with silica column chromatography. These findings reveal the antifungal and possible commercial potential of the plant extract and its potential in agriculture against pest management and food spoilage.

Aim: To determine the antifungal activity of *Piper betle* L. extracts.

Keywords: Antifungal; Ayurveda; Fourier transform infra-red spectroscopy; Nuclear magnetic resonance spectroscopy; Food spoilage; Agriculture.

Introduction

In recent years, the increase in resistance in known fungal pathogens to the available antifungal drugs has raised enormous challenges to public health issues [1-3]. In addition, conventional antifungal drugs have undesirable side effects and are very toxic such as chlorhexidine, imidazole and amphotericin B [4]. One important demand of critical significance in this context is to search for novel antifungal agents that would be less toxic and more effective. Interestingly; several medicinal plants have been extensively investigated in order to find novel bioactive compounds [5]. Moreover, several studies have suggested that a number of plant species possesses promising antimicrobial

compounds [6-9]. *Piper betle* Linn. (*Piperaceae*), a slender creeping plant, is widely distributed in India, Sri Lanka, Thailand and other tropical countries. This plant has deep green heart shaped, smooth, shining and long stalked leaves, with pointed apex. Betel leaf possess strong aromatic flavor and have been long in use for the preparation of traditional Indian ayurvedic herbal remedies. It has been reported for the treatment of various diseases such as conjunctivitis, boils and abscesses, cuts and injuries etc. [5,10]. In addition, it also acts as a breath freshener, a digestive and pancreatic lipase stimulant and a pain killer in joint pain [11,13]. Even though all these positive effects of betel leaf are known, the biochemicals of these favorable effects remain obscure. Moreover, betel leaf extract has previously been shown to have strong antimicrobial effects in review [14-19]. A couple of research articles [20,21] have demonstrated the potential of the leaf extract on dermatophytes. Here in, we show that the leaf extract possess antifungal activity against various plant pathogens. The present study was sought to investigate the effects of ethyl acetate, hexane and ethanol-methanol extracts of this plant leave on fungal pathogens.

Here in, we employed biophysical techniques to identify the active metabolite that provide the betel leaf extract with its anti-fungal and other pharmacological properties that have been valued in traditional Indian ayurveda.

Materials and methods

Betel leaf extract

The crude extract and isolation was done using Silica Gel (100-200 mesh) for column chromatography and HPLC grade solvents. Betel leaf was collected from Western Maharashtra farm (Indapur tehsil), India. It was cleaned first with distilled water and dried in shadow for a week. The dried leaves were powdered and 100 g was used for extraction. The powder was transferred into 1 L conical flask with 500 mL ethanol to completely soak the powder, which was incubated at room temperature for 24 hours.

The sample was filtered using ordinary filter paper directly into the clean round bottom flask and set for distillation. We obtained up to 30 mL of distillate from a single run under controlled heating at 55 ° C under reduced pressure. We repeated the procedure 3 times and pulled the collected distillate containing crude extract. The crude extract was concentrated on a water bath at 55 ° C under reduced pressure.

During the extraction optimization step, we added 100 mL ethyl acetate/ethanol-methanol/ hexane to the concentrated 100 mL crude extract and mixed well. The mixture was separate during separating funnel and the ethyl acetate/n-hexane fraction was collected in another conical flask. All 3 extracts were concentrated in procedure mentioned above and was further purified using column chromatography using the respective solvent .The eluent was pooled and analyzed by thin-layered chromatography (TLC), Silica gel 60 F254, preloaded Silica gel on Alumina sheets with ethyl acetate as the solvent and observed under UV chamber and Iodine vapors.

Fungal strains and growth conditions

We tested the effect of betel leaf extract on opportunistic fungal pathogen *Aspergillus niger*, which is the causative agent of black mold in several fruits and vegetables [22]. We also isolated saprophytic and opportunistic pathogenic fungi *Rhizopus sp.* [23] from the leaves of *Murraya koenigii*. (curry leaves). Lastly, we isolated a mold from the leaf of figs infested with rust, which was not obligate parasite *Cerotelium (Physopella) fici* [24] but we have tentatively identified it as wild *Aspergillus sp.* These cultures were cultured and maintained on Potato Dextrose Agar (PDA- HIMEDIA) at 4 ° C. A stock inoculums spore suspension of each fungal culture was prepared from fresh, mature (3-daysold) cultures grown on potato dextrose agar plates at 28 ° C.

In-vitro antifungal assay

Antifungal activity of betel leaf extract was tested against *A.niger*, wild *Aspergillus sp.* and *Rhizopus sp.* of the ethyl acetate, hexane and ethanol-methanol extracts of betel leaf sample was tested by well diffusion method. In brief, 500 µL of fungal spore suspension was added to 20 mL PDA medium and poured in petri dish. After solidification, wells of 5mm in diameter were made on this plate. Each well was filled with 50 µL of ethyl acetate, hexane and ethanol-methanol herbal extract. Potassium tellurite was used as positive control and ethyl acetate, hexane and ethanol-methanol solvent used as negative control. The antifungal assay plates were incubated at 30 ° C for 36h. The antifungal activities of the extracts were determined by measuring the diameter of the inhibition zone in millimetres (mm).

NMR and FTIR

The IR spectra of neat sample were recorded on Nicolette iD5, Thermo scientific at room temperature. Standard¹ H NMR spectra was recorded on Jeol 200MHz using CDCl₃ solvent and TMS (Euriso-top) as a reference at room temperature.

Results

Isolation of active ingredient from *Piper betle* extract

The beetle leaf extract was purified to homogeneity. During the solvent extraction steps, we observed three spots on the TLC plates. After solvent extraction the crude extract was concentrated and run on a column containing 100-200 mesh Silica Gel and Sodium Sulfate for further purification. In all three solvent types that we tested we observed a single spot from there fraction.

Antifungal susceptibility assay

In the next step, we studied the effect of the organic extract against a range of bacterial and fungal cultures. In all tested cases, we observed antimicrobial effect of the extract against fungal cultures only. This is in stark contrast to preciously reported data where few groups have observed anti-bacterial effect of the *Piper betle* extract [14,17,25,26]. Results obtained in the present study revealed that the ethyl acetate extract possess effective antifungal activity against all the tested fungal cultures (Table-1, Figure-1).The highest antifungal activity of ethyl acetate extract was observed for *A.niger* and Black rust followed by *Rhizopus sp.* Antifungal activity of hexane extract was also significant against *A.niger* (5 mm) and *Aspergillus sp.* (8 mm), while ethanol-methanol extract was ineffective against any of the tested fungal cultures. In addition to the molds, we also tested the effects of wild yeasts that were isolated from leaves. We again noticed a strong anti-fungal activity against them. From all the preparations, ethyl acetate fraction provided the best results. This could be because the extraction process is more efficient and perhaps also stable over other solvent.

Biophysical characterization of the extract

We performed NMR and FTIR spectroscopy in order to understand the functional groups associated with the isolated active compound. In the ¹H-NMR spectrum, we observed three

Fungal culture	Inhibition zone (mm)			
	Ethyl acetate	Hexane	Ethanol-Methanol	K-tellurite (+ control)
<i>Aspergillus niger</i>	28	5	-	6
<i>Aspergillus sp.</i>	28	8	-	7
<i>Rhizopus sp.</i>	23	-	-	-

Antifungal activity of ethyl acetate, hexane and ethanol-methanol herbal extract against indicated fungi.




Figure 1: Antifungal activity of ethyl acetate, hexane and ethanol-methanol extracts of herbal sample against A) *A.niger* B) *Aspergillus sp.* and C) *Rhizopus*. Extract preparation in a) Hexane, b) Ethanol-Methanol, c) Ethyl acetate and positive control d) Potassium tellurite

types downfield peaks at 6.7, 7.2 and 7.25 ppm for aromatic ring. Additionally plenty of peaks were observed for protons associated with polar and aliphatic groups in the 3.25-5.5 and 1-2.25 ppm, respectively. From the characteristics of the spectra and the results published by other groups, the extracted compound is a derivative of the phenyl propanoid family to which antimicrobials eugenol-chavicol belong [27]. The FTIR spectra well corroborated with the NMR spectrum. We observed a broad peak in the range of 3700 to 3000, which corresponds to absorption caused by N-H, C-H and O-H single bonds. We did not observe any peaks characteristic of triple bonds in the range of 2,500 to 2,000. A wide range of double bond specific groups in C=O and C=C was observed in the region from 2,000 to 1,500. Lastly, the region from 1700-600 show finger print that is reminiscent of phenyl propanoid eugenol [27,28].

Discussion

The antifungal activity of the various extract highlights that ethyl acetate based extraction process provided the best for antifungal property. Surprisingly, unlike other results where *Piperbetle* extract that showed both anti- bacterial and fungal activities, we did not observe any anti-bacterial effect against gram positive *Staphylococcus aureus* and *Bacillus sp.*; few gram negative bacteria such as *Escherichia coli*, *Pseudomonas aeruginosa* and *Xanthomonas campestris*. This may be due to the fact that our purification method is different to the ones reported before (Table 2). In all previous studies crude extracts have been used hence many groups have reported anti-bacterial, anti-fungal and in some cases anti-cancer effects as well. In our methodology, we have enriched the phenyl propanoid derivative in the extract by first ethanolic extract and then treating with other organic solvents like ethyl acetate/ hexane/




Figure 2: 1D ¹H-NMR spectrum of the active ingredient of *Piper betle* extract is suggestive of molecule from the phenyl propanoid- eugenol family




Figure 3: FTIR spectrum of the active ingredient of *Piper betle* extract.

Active compound	Extraction method	Result
Hydroxychavicol	choloroform extract	anti-fungal (Ali et al., 2010); antibacterial (Sharma et al 2009)
	hydrodistillation	anti-bacterial (Basak and Guha, 2015)
	Concoction- boiling extraction	anti-cancer (Gundala et al., 2014)
Eugenol	ethanol or water-soxhlet extraction	<i>Streptococcus mutans</i> (Deshpande and Kadam, 2013)
	solvent extraction: methanol, ethyl acetate and petroleum ether	<i>Streptococcus mutans</i> (Deshpande and Kadam, 2015)
	hydro-distillation	anti-bacterial (Sugumaran et al., 2011)
Hydroxychavicol and Eugenol	methanolic extract	anti-cancer (Paranjpe et al., 2013)
	liquid-liquid and supercritical fluid extraction	Comparison of extraction methods(Singtongratana et al., 2013)

Summary of bio-active extraction procedures from *Piper betle*

ethanol-methanol. Furthermore, the large zone of clearance that we observe highlights the better extraction and stability of the phenyl propanoid in the ethyl acetate fraction.

Conclusion

Our finding sheds light on one of the important biochemical metabolite that contributes to the significance of betel leaf in traditional Indian ayurvedic medicine. The metabolite can be readily extracted in a two-step process using organic solvents and silica column chromatography. The active ingredient belongs to the phenyl propanoid family belonging to eugenol-chavicol group have been shown to possess strong antimicrobial properties [15,26,29,30]. More importantly, our results highlight the potential of using betel leaf extract as a potent anti-fungal agent for farming and perhaps also food storage against different types of molds. Food spoilage is a major agricultural problem accounting for heavy losses; therefore it is necessary to make the process commercially viable and efficacious against various pathogens and food spoilage organisms [31,32].

Acknowledgements

We thank the following institutes like M. E. S Abasaheb Garware college, Vijay Chemicals and Neeti Developer for support; and Dr. Nivedita Cukkemane for critically reading the manuscript.

References

- Alexander BD, Perfect JR, Antifungal resistance trends towards the year 2000. Implications for therapy and new approaches. *Drugs*. 1997;54(5):657-678.
- Ghannoum MA, Rice LB, Antifungal agents: mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance. *Clin Microbiol Rev*. 1999;12(4):501-517.
- Neely MN, Ghannoum MA. The exciting future of antifungal therapy.

- Eur J Clin Microbiol Infect Dis. 2000; 19(12):897–914.
4. White TC, Marr KA, Bowden RA, 1998. Clinical, cellular and molecular factors that contribute to antifungal drug resistance. Clin Microbiol. 1998;11(2):382–402.
 5. Peter KV. Handbook of herbs and spices. 1st Edition Vol 1 and Vol 2. Sawston, Cambridge: Woodhead Publishing Limited;2004.
 6. Nakamura CV, Ishida K, Faccin LC, Filho BPD, Cortez DAG, Rozenal S, et al. In vitro activity of essential oil from *Ocimum gratissimum* L. against four *Candida* species. Res Microbiol. 2004;155(7):579–586. doi:10.1016/j.resmic.2004.04.004
 7. Wannissorn B, Jarikasem S, Siriwangchai T, Thubthimthed S. Antibacterial properties of essential oils from Thai medicinal plants. Fitoterapia. 2005;76(2):233–236.
 8. Khan R, Islam B, Akram M, Shakil S, Ahmad A, Ali SM, et al. Antimicrobial Activity of Five Herbal Extracts Against Multi Drug Resistant (MDR) Strains of Bacteria and Fungus of Clinical Origin. Molecules. 2009;14(2):586–597. doi: 10.3390/molecules14020586
 9. Jagtap SD, Deokule SS, Pawar PK, Harsulkar AM, Kuvalekar A A. Antimicrobial activity of some crude herbal drugs used for skin diseases by *Pawra* tribes of Nandurbar district. J Nat Prod. 2010;1(2):216–220.
 10. Guha P. Betel leaf: The Neglected Green Gold of India. J Hum Ecol. 2006;19(2):87–93.
 11. Norton SA. Betle: consumption and consequences. J Am Acad Dermatol. 1998;38(1):81–88.
 12. Prabhu MS, Patel K, SaraawathiG, Srinivasan K, Effect of orally administered betel leaf (*Piperbetel leaf* Linn.) on digestive enzymes of pancreas and intestinal mucosa and on bile production in rats. Indian J Exp Biol. 1995;33(10):752–756.
 13. Pradhan D, Suri KA, Pradhan DK, Biswasroy P. Golden Heart of the Nature: *Piper betle* L. Journal of Pharmacognosy and Phytochemistry. 2013;1(6).
 14. Ramji N, Ramji N, IyerR, Chandrasekaran S, 2002. Phenolic antibacterial from *Piper betle* in the prevention of halitosis. J Ethnopharmacol. 2002;83(1-2):149–152.
 15. Sugumaran M, Suresh Gandhi M, Sankarnarayanan K, Yokesh M, Poornima M, Sree Rama rajasekhar, Chemical composition and antimicrobial activity of vellaikodivariety of *Piper betle* Linn Leaf oil against dental pathogens. Int J PharmTech Res. 2011;3(4):2135–2139.
 16. Datta A, Ghoshdastidar S, Singh M. Antimicrobial Property of *Piper betle* leaf against Clinical Isolates of Bacteria. IJPSR. 2011;2(3):104–109.
 17. Kaveti B, Tan L, Antibacterial Activity of *Piper betle* Leaves. International Journal of Pharmacy Teaching and Practices. 2011;2(3):129–132.
 18. Chakraborty D, Shah B. Antimicrobial, anti-oxidative and anti-hemolytic activity of *Piper betel leaf* extracts. IJPPS. 2011;3(3):192–199.
 19. Das S, Parida R, Sriram Sandeep I, Nayak S, Mohanty S. Biotechnological intervention in betelvine (*Piper betle* L.): A review on recent advances and future prospects. Asian Pac J Trop Med. 2016;9(10):938–946. doi:10.1016/j.apjtm.2016.07.029
 20. Trakranrungsie N, Chatchawanchonteera A, Khunkitti W. Ethnoveterinary study for anti-dermatophytic activity of *Piper betle*, *Alpinia galangal* and *Allium ascalonicum* extracts *in vitro*. Res Vet Sci. 2008;84(1):80–84.
 21. Sharma KK, Saikia R, Kotoky J, Kalita JC, Das J. Evaluation of Anti-dermatophytic activity of *Piper betle*, *Allamanda cathertica* and their combination: An *in vitro* and *in vivo* study. Int J PharmTech Res. 2011;3(2):644–651.
 22. Sharma R. Pathogenicity of *Aspergillus niger* in plants. Cibtech Journal of Microbiology. 2012;1(1):47–51.
 23. Wilson CL, Wisniewski ME. Biological control of postharvest diseases of fruits and vegetables: an emergency technology. Annual Review of Phytopathology. 1989;27:425–441. doi: 10.1146/annurev.py.27.090189.002233
 24. Laundon GF, Rainbow AF. *Cerotelium fici*. CMI Descriptions of Pathogenic Fungi and Bacteria. 1971;281:1–2.
 25. Deshpande SN, Kadam DG. GCMS analysis and antimicrobial activity of Piper betle (Linn) leaves against *Streptococcus mutans*. Asian J Pharm Clin Res. 2013;6(5):99–101.
 26. Basak S, Guha P. Modelling the effect of essential oil of betel leaf (*Piper betle* L.) on germination, growth, and apparent lag time of *Penicillium expansum* on semi-synthetic media. Int J Food Microbiol. 2015;215:171–178. doi:10.1016/j.ijfoodmicro.2015.09.019
 27. Carrasco AH, Espinoza CL, Cardile V, Gallardo C, Cardona W, Lombardo L, Catalán MK, Cuellar FM, Russo A. Eugenol and its synthetic analogues inhibit cell growth of human cancer cells (Part I). Journal of the Brazilian Chemical Society. 2008;19:543–548.
 28. Paranjpe R, Gundala SR, Lakshminarayana N, Sagwal A, Asif G, Pandey A, et al. Piper betel leaf extract: anticancer benefits and bio-guided fractionation to identify active principles for prostate cancer management. Carcinogenesis. 2013;34(7):1558–1566. doi: 10.1093/carcin/bgt066
 29. Prakash B, Shukla R, Singh P, Kumar A, Mishra PK, Dubey NK. Efficacy of chemically characterized *Piper betle* L. essential oil against fungal and aflatoxin contamination of some edible commodities and its antioxidant activity. Int J Food Microbiol. 2010;142(1-2):114–119.
 30. Saxena M, Khare NK, Saxena P, Syamsundar KV, Srivastava SK. Antimicrobial activity and chemical composition of leaf oil in two varieties of *Piper betle* from northern plains of India. JSIR. 2014;73(2):95–99.
 31. Rawat S. Food Spoilage: Microorganisms and their prevention. Asian J. Plant Sci. Res. 2015;5(4):47–56.
 32. Adeyeye SAO. Fungal mycotoxins in foods: A review. Cogent Food & Agriculture. 2016;2(1). doi: 10.1080/23311932.2016.1213127
 33. Ali I, Khan FG, Suri KA, Gupta BD, Satti NK, Dutt P, et al. In vitro antifungal activity of hydroxychavicol isolated from *Piper betle* L. Annals of Clinical Microbiology and Antimicrobials. 2010;9:1–7. doi: 10.1186/1476-0711-9-7
 34. Sharma S, Khan IA, Ali I, Ali F, Kumar M, Kumar A. et al. Evaluation of the Antimicrobial, Antioxidant, and Anti-Inflammatory Activities of Hydroxychavicol for Its Potential Use as an Oral Care Agent. Antimicrobial Agents and Chemotherapy. 2009;53(1):216–222. doi:10.1128/AAC.00045-08
 35. Gundala SR, Yang C, Mukkavilli R, Paranjpe R, Brahmabhatt M, Pannu V. et al. Hydroxychavicol, a betel leaf component, inhibits prostate cancer through ROS-driven DNA damage and apoptosis. Toxicol Appl Pharmacol. 2014;280(1):86–96. doi: 10.1016/j.taap.2014.07.012
 36. Deshpande SN, Kadam DG. Evaluation of antimicrobial activity of extracts of Piper betle (Linn) leaves against *Streptococcus mutans*. World Journal of Pharmacy and Pharmaceutical Sciences. 2015;4(11):1040–1050.
 37. Singtongratana N, Vadhanasin S, Singkhonrat J. Hydroxychavicol and Eugenol Profiling of Betel Leaves from *Piper betle* L. Obtained by Liquid-Liquid Extraction and Supercritical Fluid Extraction. Kasetsart J (Nat Sci). 2013;47:614–623.
 38. Naik PM, Al-Khayri JM. Impact of Abiotic Elicitors on In vitro Production of Plant Secondary Metabolites: A Review. J Adv Res Biotech. 2016;1(2):7.