Chitinase and Glucanase Activities of Antagonistic *Streptomyces* Spp Isolated From Fired Plots under Shifting Cultivation In Northeast India

Mukesh K Malviya¹, Pankaj Trivedi² and Anita Pandey^{3*}

¹Crop Genetics Improvement and Biotechnology Lab, Guangxi Academy of Agricultural Sciences, Nanning, 530007, China

²Department of Bioagricultural Sciences and Pest Management, Colorado State University, C034 Plant Sciences, Fort Collins - CO - 80523-1177

³Biotechnological Applications, G.B. Pant National Institute of Himalayan Environment and Sustainable Development, Kosi-Katarmal, Almora, 263 643 Uttarakhand, India

Received: 10 November, 2017; Accepted: 19 January, 2018; Published: 31 May 2018

***Corresponding author:** Anita Pandey, Biotechnological Applications, G. B. Pant National Institute of Himalayan Environment and Sustainable Development, Kosi-Katarmal, Almora, 263 643 Uttarakhand, India; E-mail: anita@gbpihed.nic.in

Abstract

Antagonistic *Streptomyces* spp (*Streptomyces* sp NEA55 and *Streptomyces cavourensis* NEA5), isolated from fired plots under shifting cultivation in northeast India, are studied for their chitinase and glucanase activities. The species showed strong antagonism against test fungi (*Rhizoctonia solani* and *Cladosporium* sp.) in plate assays. Maximum % inhibition was observed due to the effect of diffusible compounds produced by these species. *Streptomyces* sp. NEA55 showed 54.83 % inhibition against *R. solani* while *S. cavourensis* NEA5 showed up to 50.00 % inhibition against *Cladosporium* sp. The inhibitory effect of volatile compounds by *Streptomyces* sp NEA55 was recorded up to 50.7 % against *R. solani* and 37.50 % against *Cladosporium* sp. While *S. cavourensis* NEA5 showed 49.23 % inhibition against *R. solani* and 34.37 % inhibition against *Cladosporium* sp. *S. cavourensis* NEA5 and *Streptomyces* sp. NEA55 produced 0.138 ± 0.006 $\mu\text{g/ml}$ and 0.15 ± 0.004 $\mu\text{g/ml}$ chitinase, 0.22 ± 0.001 $\mu\text{g/ml}$ and 0.25 ± 0.002 $\mu\text{g/ml}$ β 1,3 glucanase, respectively. Both the species showed maximum chitinase activity at pH 6 and temperature 50 °C, while minimum enzyme activity was observed at pH 10 and temperature 20 °C. Both the species showed glucanase activity maximum at pH 7 and temperature 40 °C and minimum activity at pH 10 and temperature 20 °C. Both the species hydrolyzed glycol-chitin as a substrate in denaturing conditions showing variable amount of different isoforms. This study demonstrates that the antagonistic species of *Streptomyces* survive the fire operations under shifting cultivation.

Keywords: *Streptomyces*; Antagonism; Chitinase; Glucanase; Shifting cultivation

Introduction

Actinobacteria are most widely distributed and distinct group of microorganisms in nature. *Actinobacteria*, *Streptomyces* species in particular, has been a broadly exploited group of microorganisms for the production of important secondary metabolites and enzymes in the field of medicine and agriculture

[1, 2]. *Streptomyces* are well known as antifungal biocontrol agents that inhibit several plant pathogenic fungi [3]. The antagonistic activity of *Streptomyces* to fungal pathogens is usually related to the production of antifungal compounds and extracellular hydrolytic enzymes [4, 5]. Many species of *Streptomyces* are well known as antifungal biocontrol agents that inhibit several plant pathogenic fungi e.g., *Phytophthora capsici*, *Sclerotinia rolfii*, *Fusarium sporotrichiodes*, *Rhizoctonia solani* and *Sclerotium rolfii*, *Alternaria alternata* and *Phomopsis archeri* [6-9]. Furthermore, *Streptomyces* produce bioactive compounds such as antimicrobial, antiparasitic and immune-suppressing compounds via secondary metabolism. *Streptomyces* have been found in beneficial associations with plants where they improve plant growth and protect against pests; this has attracted the attention of researchers worldwide [10].

Chitinase and β -1,3-glucanase are considered to be important hydrolytic enzymes in the lysis of fungal cell walls [11]. Chitinolytic enzymes have been identified in several *Streptomyces* spp. including *Streptomyces* sp. M-20, *S. venezuelae* P10, and *S. anulatus* CS242 [12-14]. Glucanase has been known to be produced by several microorganisms and playing important role in biocontrol [15]. Several *Streptomyces* have been studied for antifungal properties along with the production of glucanase, some of the examples are *Streptomyces* sp. S27 and *Streptomyces* sp. Mo [16, 17]. Shifting cultivation, refers to 'slash and burn', is a predominant form of agricultural practice in hills of northeast India. The microbiological aspects, basically survival of bacterial, fungal and actinobacterial communities after fire events have been studied in recent times [18-20]. The focus of present study is on the antagonistic potential of two *Streptomyces* species that were isolated after the fire events. These species have been studied with respect to production of diffusible and volatile compounds

against test fungi along with the chitinase and β -1, 3-glucanase activities. In addition, both the species are also studied with hydrolyzed glycol-chitin as a substrate in denaturing conditions.

Material and Methods

Study site and isolation of actinobacteria

Actinobacteria were isolated from the soil samples that were collected after the completion of fire events in Papumpare District, Itanagar, Arunachal Pradesh under shifting cultivation. The details of the study sites have been described in Pandey et al. [18]. Among the two actinobacterial isolates used in the present study, isolate no. NEA5 showed maximum similarity with *Streptomyces cavourensis* NR_043851 and NEA55 with *Streptomyces* sp. YIM8 AF389344 [19]. The test fungi (*Rhizoctonia solani* and *Cladosporium* sp.) were also originally isolated from shifting cultivation site [20].

Scanning electron microscopy

In order to see the deep morphological pattern (substrate and aerial mycelia) of *Streptomyces* spp. scanning electron microscopy was performed. Glutaraldehyde (2.5 %) was added to the culture and then centrifuged. After centrifugation, washing was given with phosphate buffer saline (PBS) twice and centrifuged. The sample was dehydrated with CPD (Critical dry point), and then the samples were coated with gold by auto fine coater (JFC-1600). After coating, the sample was viewed under scanning electron microscope (JSM-6610LV).

Dual culture technique for determination of the production of diffusible antifungal compounds

To test the ability of the *Streptomyces* spp. to inhibit the phytopathogens, the test fungal culture and *Streptomyces* sp. was spot inoculated off-center on each potato carrot agar (PCA) plate. After 7 days of incubation at 28 °C, the zone of inhibition was measured. Per cent growth inhibition was calculated using the following formula: $(R1-R2)/R1 \times 100$ (where R1 represents the radius of the test fungus in the direction with no bacterial colony and R2 is the radius of the fungal colony in the direction of the bacterial colony) as described in Trivedi et al. [21].

Sealed double Petri plate technique for determination of the production of volatile antifungal compounds

Each *Streptomyces* sp. was streaked on Petri plate containing PCA and a 6.0 mm disc of 4 days old culture of the test fungus was placed in the middle of another PCA plate. The lid of both the plates were replaced with the plates of same diameter, placed face to face, and sealed by parafilm, preventing any physical contact between the pathogen and the *Streptomyces* sp. Petri plates were incubated at 28 °C for 7 days and growth of the pathogen was measured and compared to control developed in the absence of the antagonist (mocked inoculation with a 6.0 mm disc of PCA). Results were expressed as means of per cent inhibition of the growth of each fungus in the presence or absence of the antagonistic *Streptomyces* sp. Per cent growth inhibition was

calculated by the following formula: $(R1-R2)/R1 \times 100$, where R1 represents the diameter of the test fungus on control plate and R2 is the diameter of the growth on inoculated plate.

Quantitative estimation of production of chitinase and glucanase on different temperature and pH

Quantification of chitinase and glucanase has been done following prescribed procedures [22]. The *Streptomyces* species were cultured at 28 °C for 5 days on a rotary shaker in 250.0 ml of chitin peptone medium for chitinase production and peptone medium containing laminarin (0.2 %) (from *Laminaria digitata*, Sigma) for glucanase production [23]. The cultures were centrifuged at 12,000 g for 20 min at 4 °C and the supernatants were used as enzyme source. The reaction mixture contained 0.25 ml of enzyme solution, 0.3 ml of pH buffer (1.0 M citrate buffer pH 5, phosphate buffer pH 6 to 8, and glycine buffer pH 9). The reaction mixtures were incubated at different temperatures (20, 30, 40, 50 and 60 °C) for 4 h in a water bath. One unit of chitinase was determined as 1 μ mol of N acetyl glucosamine (GlcNAc) released min⁻¹ and β -1, 3-glucanase activity was determined as 1 μ mol of glucose released min⁻¹. Protein content was determined as described by Lowry method [24].

SDS-PAGE analysis for chitinase

Streptomyces spp. was grown as shake culture in chitin-peptone medium at 28 °C for 7 days. The supernatant was centrifuged at 12000 g for 20 min at 4 °C and filtered through 0.22 μ l sterile filter (Millipore) and collected in conical flask. Protein content was analyzed by SDS-PAGE [25]. Polyacrylamide slab gel consisted of 4 % stacking gel and 10 % separating gel containing 0.01 % glycol-chitin. Electrophoresis was carried out at a constant voltage of 65 V; gel was stained with coomassie brilliant blue R-250 and analyzed under gel documentation system (Alpha Imager 2200). Glycol chitin was prepared as described by Trudel and Asselin [26].

Results and Discussion

The scanning electron microscopy of both the species is presented in Figure: 1A and Figure: 1B. Scanning electron microscopy revealed definite structures of mycelium and spores of both the *Streptomyces* spp. Two of the isolates that were used for detailed studies based on their antagonistic properties were subjected to scanning electron microscopy. The scanning electron microscopy has been referred to provide perfect characterization of *streptomyces* [27]. The effect of diffusible compounds produced by the *Streptomyces* spp., evaluated in terms of reduction in radial growth of two test fungi viz. *Rhizoctonia solani* and *Cladosporium* sp., following 7 days of incubation at 28 °C, are presented in Figure:2 & Table 1. The results showed that volatile compounds produced by *Streptomyces* spp. were inhibitory to the growth of the test fungi, viz. *R. solani* and *Cladosporium* sp. Inhibition (%) in fungal growth by *Streptomyces* species is presented in Table 1. Microscopic observations revealed that the diffusible as well as volatile compounds produced by

Figure 2: Production of antifungal compounds by *Streptomyces cavourensis* NEA5 & *Streptomyces* sp. NEA55 (A&B) respectively: Inhibition of *R. solani* and *Cladosporium* sp. due to diffusible compounds produced. Morphological deformities: (A1&B1) Normal structures of *R. solani* and *Cladosporium* sp. (A2&B2) deformed structures of respective fungus. (Bar = 5 µm)

Table 1: In vitro percent inhibitory effect of diffusible and volatile metabolites of <i>Streptomyces</i> species on growth of test fungi				
Pathogenic test fungi	% Inhibition in fungal growth by <i>Streptomyces</i> spp. after 7 days of incubation			
	<i>Streptomyces cavourensis</i> NEA5		<i>Streptomyces</i> sp NEA55	
<i>Rhizoctonia solani</i>	51.61	49.23	54.83	50.07
<i>Cladosporium</i> sp.	50	34.37	47.7	37.5

Table 2: Comparative account of structures of normal and antagonized pathogen

Test fungi	Structure	Normal	Antagonized
<i>Rhizoctonia</i>	Mycelium	Somatic and fertile hypha well distinguished	Irregular
<i>Solani</i>	Somatic hypha	Septate, dia. 2.5- 3.5 μ m	Irregular
	Fertile hypha	Septate and dark in colour, dia. 3.0-4.5 μ m	Septate, dia 3.0-6.0
	Hyphal wall	Present	Irregular septate, dia 3.0-7.5 μ m
<i>Cladosporium</i> sp.	Mycelium	Somatic hypha septate, dia. 2-3.5 μ m	Lysed
	Conidia	Curved or straight width 2.0-2.5 μ m	Somatic hypha septate, dia. 2-5 μ m
			Irregular width 2-4 μ m

Streptomyces species induced morphological abnormalities in the fungal structures (Table 2). Deformation was observed in *mycelial*, *hyphal* or *conidial* structures. The longitudinal septae completely disappeared and the conidia became thick walled and spherical or irregular in shape. Lysis of fungal hyphae and vacuolization as well as granulation in mycelium was observed in *R. solani*. Size of the somatic and fertile hypha of *R. solani* increased due to the antagonistic effect of actinobacteria. Due to the antifungal activity of *Streptomyces* sp., hyphal wall of *R. solani* was completely lysed. In case of *Cladosporium* sp., size of mycelium and conidia increased with irregular shape of conidia. The inhibition of growth of the test pathogenic fungi continued to increase with increasing incubation time. A clear inhibitory area and dense sporulation ring was observed near the growth of the isolate which antagonized respective test fungi. No physical contact was observed between the *Streptomyces* species and the test fungi that were observed to be antagonized. Moreover, the formation of the inhibitory halo suggested the presence of fungicidal metabolites secreted by *Streptomyces*. Similar observations have been reported by Aghigni et al. [28] from a number of actinobacterial isolates from Iranian soil. These isolates formed inhibition zones in dual culture based assays inhibiting the growth of *Alternaria solani*, *A. alternata*, *Fusarium solani*, *Phytophthora megasperma*, *Verticillium dahlia* and *Saccharomyces cerevisiae*. The antifungal potential of actinobacteria against *Colletotrichum gloeosporioides* and *Sclerotium rolfsii* was assessed by dual culture technique [10]. Al-Askar et al. [29] reported that *Streptomyces spororaveus* RDS28 produces antifungal compounds against some phytopathogenic fungi, viz., *Rhizoctonia solani*, *Fusarium solani*, *F. verticillioides*, *Alternaria alternata* and *Botrytis cinerea*. There was a change in colour of the test fungi in inoculated plates, indicative of the inhibitory effect of volatile compounds. The test fungi, inhibited by the *Streptomyces*, under present study exhibited morphological abnormalities due to the production of diffusible and volatile antifungal compounds. Abnormal hyphal swelling, degradation and lysis of mycelia were observed by Joo [6] when *Phytophthora capsici* was grown with the high or low molecular fraction of *Streptomyces halstedii* AJ-7.

Quantitative estimation of production of chitinase and β 1, 3 glucanase activities is presented in Figure 3 (A&B) showing maximum activity of chitinase and glucanase. Effect of different

pH and temperature on chitinase activity of both the species is presented in Figure 4 (A&B). Both the species showed maximum activity of chitinase at pH 6 and temperature 50 °C, while minimum enzyme activity was observed at pH 10 and temperature 20 °C. In case of glucanase activity both the species showed maximum enzyme activity at pH 7 and temperature 40 °C and minimum activity at pH 10 and temperature 20 °C (Figure 5A&B). Kim et al. [12] reported maximum activity of chitinase

Figure 3A: Quantitative estimation of chitinase activity

Figure 3B: Quantitative estimation of glucanase activity

Figure 4A: Chitinase activity of *Streptomyces cavourensis* NEA5 at different pH & temperature

Figure 4B: Chitinase activity of *Streptomyces* sp NEA55 at different pH & temperature

Figure 5A: Glucanase activity of *Streptomyces cavourensis* NEA5 at different pH & temperature

between pH 5.0 to 6.0 and it was relatively stable at pH 4.0 to 8.0 when kept at 4 °C. Beyond this pH range there was a rapid loss in the activity. Glucanase activity of the two isolates was found maximum at pH 7 and temperature 40 °C. Highest activity was found between pH 5 to 8 and temperature 30 to 50 °C, in previous studies. The optimum activity for short term incubation is often seen at temperature in the range of 30 to 50 °C, while many fungal glucanases appear stable between 50 to 60 °C [30].

SDS-PAGE exhibited the chitinase activity of the two *Streptomyces* species (NEA5 & NEA55). *Streptomyces* species hydrolyzed glycol-chitin as a substrate in denaturing conditions. Species showing variable amount of different isoforms with molecular weights between 31 to 40 kDa are presented in Figure 6. Majority of bacterial chitinases has been reported to be in the range of 20 to 60 kDa. Chitinases from various *Streptomyces* were found to possess molecular weights as 20 kDa from *Streptomyces*

Figure 5B: Glucanase activity of *Streptomyces* sp NEA55 at different pH & temperature

Figure 6: Detection of chitinase activity after SDS-PAGE using glycol chitin as substrate, lane 1 (Molecular weight markers), lane 2 and 3 sample *Streptomyces cavourensis* NEA5 and *Streptomyces* sp NEA55, respectively

sp. M-20 [13] and some strains of *Streptomyces* varying from approximately 25 kDa to 200 kDa [31].

The aim of this work is to prove the potential of *Streptomyces* species to survive the fire operations retaining their potential to reduce or eliminate the plant pathogenic fungi. Such robust isolates may prove to be the potential biocontrol agents for field applications. Further studies are needed in order to determinate the nature of *Streptomyces* species metabolites and their mechanism of action.

Acknowledgment

Director (GBPNIHESD) for extending the facilities and Ministry of Environment, Forest and Climate Change, Govt. of India, New Delhi for financial support.

References

- Kumar D, Gupta RK. Biocontrol of wood rotting fungi. *Ind J Biotechnol.* 2006; 5:20-25.
- Narayana KJP, Vijayalakshmi M. Chitinase production by *streptomyces* sp. anu 6277. *Braz J Microbiol.* 2009; 40(4): 725-733.
- El-Tarabily KA, Soliman MH, Nassar AH, Al-Hassani HA, Sivasithamparam K. Biocontrol of sclerotinia minor using a chitinolytic bacterium and actinomycetes. *Plant Pathol.* 2000; 49: 573-583.
- Fourati-Ben Fguira L, Fotso S, Ben Ameer-Mehdi R, Mellouli L, Laatsch H. Purification and structure elucidation of antifungal and antibacterial activities of newly isolated *Streptomyces* sp. strain US80. *Res Microbiol.* 2005; 156(3): 341-347.
- Trejo-Estrada SR, Sepulveda IR, Crawford DL. In vitro and in vivo antagonism of *Streptomyces violaceusniger* YCED9 against fungal pathogens of turfgrass. *World J Microbiol Biotechnol.* 1998;14(6):865-872.
- Joo GJ. Production of an anti-fungal substance for biological control of *Phytophthora capsici* causing blight in red-peppers by *Streptomyces halstedii*. *Biotechnol Lett.* 2005;27(3):201-205.
- Errakhi R, Bouteau F, Lebrihi A, Barakate M. Evidences of biological control capacities of *Streptomyces* spp. against *Sclerotium rolfsii* responsible for damping-off disease in sugar beet (*Beta vulgaris* L.). *World J Microbiol Biotechnol.* 2007;23(11):1503-1509.
- Rugthaworn P, Uraivan Dilokkunanant, Sangchote S, Piadang N, Kitpreechavanich V. Search and improvement of actinomycete strains for biological control of plant pathogens. *Kasetsart J (Nat Sci).* 2007;41:248-254.
- Malviya MK, Pandey A, Trivedi P, Gupta G, Kumar B. Chitinolytic activity of cold tolerant antagonistic species of *streptomyces* isolated from glacial sites of Indian Himalaya. *Curr Microbiol.* 2009;59(5):502-508. doi: 10.1007/s00284-009-9466-z
- Sousa JA, Olivares FL. Plant growth promotion by streptomycetes: ecophysiology, mechanisms and applications. *Chem Biol Technol Agric.* 2016; 3:24.

11. Prapagdee B, Kuekulvong C, Mongkolsuk S. Antifungal potential of extracellular metabolites produced by *Streptomyces hygroscopicus* against phytopathogenic fungi. *Int J Biol Sci*. 2008;4(5):330-337.
12. Kim KJ, Yang YJ, Kim JG. Purification and characterization of chitinase from *Streptomyces* sp. M-20. *J Biochem Mole Biol*. 2003;36(2):185-189.
13. Mukherjee G, Sen SK. Purification, characterization, and antifungal activity of chitinase from *Streptomyces venezuelae* P10. *Curr Microbiol*. 2006;53(4):265-269.
14. Mander P, Cho SS, Choi YH, Panthi S, Choi YS, Kim HM, Yoo JC et al. Purification and characterization of chitinase showing antifungal and biodegradation properties obtained from *Streptomyces anulatus* CS242. *Arch Pharm Res*. 2016;39(7):878-886. doi: 10.1007/s12272-016-0747-3
15. El-Katatny MH, Gudelj M, Robra KH, Elnaghy MA, Gubitz GM. Characterization of a chitinase and an endo- β -1,3-glucanase from *Trichoderma harzianum* Rifai T24 involved in control of the phytopathogen *Sclerotium rolfsii*. *Appl Microbiol Biotechnol*. 2001; 56(1-2):137-143.
16. Shi P, Yao G, Li N, Luo H, Bai Y, Wang Y, Yao Bin et al. Cloning, characterization, and antifungal activity of an endo-1,3- β -D: -glucanase from *Streptomyces* sp. S27. *Appl Microbiol Biotechnol*. 2010;85(5):1483-1490. doi: 10.1007/s00253-009-2187-1
17. Kurakake M, Yamanouchi Y, Kinohara K, Moriyama S. Enzymatic properties of β -1,3-glucanase from *Streptomyces* sp Mo. *J Food Sci*. 2013;78(4):502-506. doi: 10.1111/1750-3841.12076
18. Pandey A, Chaudhry S, Sharma A, Choudhary VS, Malviya MK, Chamoli S, Rinu K, Trivedi P, Palni LM et al. Recovery of *Bacillus* and *Pseudomonas* spp from the 'Fired Plots' under Shifting Cultivation in Northeast India. *Curr Microbiol*. 2011; 62(1): 273-280. doi: 10.1007/s00284-010-9702-6
19. Malviya MK, Pandey A, Sharma A, Tiwari SC. Characterization and identification of actinomycetes isolated from 'fired plots' under shifting cultivation in northeast Himalaya, India. *Ann Microbiol*. 2013;63(2):561-569.
20. Jain R, Chaudhary D, Dhakar K, Pandey A. A consortium of fungal survivors after fire operations under shifting cultivation in Northeast Himalaya, India. *Nat Acad Sci Lett*. 2016;39(5):343-346.
21. Trivedi P, Pandey A, Palni LM. In vitro evaluation of antagonistic properties of *Pseudomonas corrugata*. *Microbiol Res*. 2008;163(3):329-336.
22. Nagarajkumar M, Bhaskaran R, Velazhahan R. Involvement of secondary metabolites and extracellular lytic enzymes produced by *Pseudomonas fluorescens* in inhibition of *Rhizoctonia solani*, the rice sheath blight pathogen. *Microbiol Res*. 2006;159(1):73-81.
23. Lim H, Kim Y, Kim S. *Pseudomonas stutzeri* YLP-1 genetic transformation and antifungal mechanism against *Fusarium solani*, an agent of plant root rot. *Appl Environ Microbiol*. 1991;57(2):510-516.
24. Bollag DM, Rozycki MD, Edelstein SJ. *Protein methods*, 2nd Ed. Wiley New York. 1996:432.
25. Laemmli UK. Cleavage of structural proteins during the assembly of head of bacteriophage T4. *Nature (London)*. 1970;227:680-685.
26. Trudel J, Asselin A. Detection of chitinase activity after polyacrylamide gel electrophoresis. *Ann Biochem*. 1989; 178(2):362-366.
27. Dietz A, Mathews J. Scanning Electron Microscopy of Selected Members of the *Streptomyces hygroscopicus* Group. *Appl Microbiol*. 1969;18(4):694-696.
28. Aghigni S, Boniar GHS, Rowashdeh R, Batayneh S, Saadoun L. First report on antifungal spectra of activity of Iranian actinomycetes strains against *Alternaria solani*, *Alternaria alternata*, *Fusarium solani*, *Phytophthora megasperma*, *Verticillium dahlia* and *Saccharomyces cerevisiae*. *Asian J Plant Sci*. 2004;3(4):463-471. doi: 10.3923/ajps.2004.463.471
29. Al-Askar AA, Abdul Khair WM, Rashad YM. In vitro antifungal activity of *Streptomyces spororaveus* RDS28 against some phytopathogenic fungi. *Afr J Agric Res*. 2011;6(12):2835-2842.
30. Muskhazli M, Ramli SA, Ithni A, Tohfah NF. Purification and characterisation of β -1, 3-giucanase from *Trichodenna harzianum* BIO 10671. *Pertanika J Trop Agric Sci*. 2005; 28(1): 23-31.
31. Gomes RC, Semeˆdo LTAS, Soares RMA, Alviano CS, Linhares LF, Coelho RR et al. Chitinolytic activity of actinomycetes from a cerrado soil and their potential in biocontrol. *Lett Appl Microbiol*. 2000; 30(2):146-150.