

An Unusual Course of Necrotizing Pneumonia Leading to Pneumonectomy in A Child

Mohamed Zouari^{1,2*}, Ahmed Khalil Ben Abdallah^{1,2}, Mahdi Ben Dhaou^{1,2}, Riadh Mhiri^{1,2}

¹Department of pediatric surgery, Hedi-Chaker Hospital 3029 Sfax, Tunisia.

²Sfax Medical School, Sfax, Tunisia.

Received: February 06, 2020 ; Accepted: February 20, 2020 ; Published: March 11, 2020

***Corresponding author:** Mohamed Zouari, Department of Pediatric Surgery. Hedi Chaker Hospital. 3029 Sfax. Tunisia. Tel: +21697459586; E-mail: zouarimohamed.1982@yahoo.fr

A 13-year-old boy with a history of complicated pneumonia at the age of 20 months (Figure 1-A) managed with antibiotics and chest drainage presented with hemoptysis. Chest radiography showed multiple small, thin-walled cavities (Figure 1-B). Chest




Figure 1: PA: Chest radiograph showing whiteout of the left lung
B: Chest radiograph showing multiple small, thin-walled cavities
C, D: Axial (C) and frontal (D) CT scan cuts showing the development of multiple air-filled cavities within the lung parenchyma
E, F: Lung perfusion scan showing a reduced perfusion of the left lung

scan showed massive necrosis of the left lung (C and D). Lung perfusion scan showed differential perfusion of left: right lungs to be 8:92 % (E and F). The patient underwent left pneumonectomy. Postoperative course was uneventful and the patient did not develop respiratory tract symptoms during a 1-year follow-up period.

Necrotizing pneumonia is characterized by the development of necrosis, liquefaction, and cavitation of the pulmonary parenchyma due to pulmonary tissue lysis of infectious origin [1]. This condition is uncommon in children [2]. The major pathogens are *Staphylococcus aureus* and *Streptococcus pneumoniae* [1]. The standard treatment for necrotizing pneumonia includes empiric intravenous antimicrobial therapy with possible transition to oral antibiotics for a median duration of 28 days [3]. Lung necrosectomy should be considered at an early stage in the management of these patients, which avoids a more aggressive resection [4].