Knowledge and Attitudes of Coaches Regarding Sports Related Oro-Facial Injuries in Chennai, India

M. Priya, Ditto Sharmin*, Deepti Amarlal, Eapen Thomas and Y. Pooja

Department of Pediatric and Preventive Dentistry, Meenakshi Ammal Dental College, Alapakkam main road, Maduravoyal, Chennai

Received: May 17, 2016; Accepted: July 04, 2016; Published: July 18, 2016

***Corresponding author:** DittoSharmin, Department of Pediatric and Preventive Dentistry, Meenakshi Ammal Dental College, Alapakkam main road, Maduravoyal, 600 095, Chennai, India; Tel no: +91 9486190456; Fax- +91 44 23781631; E-mail: dr.dittosharmin@gmail.com

Abstract

Background/Aim: The aim of this study was to assess the knowledge and attitudes of sports coaches regarding sports related injuries and its prevention using a questionnaire survey design.

Material and Methods: A modified questionnaire was distributed to 50 coaches of various sports teams in different parts of Chennai city in India between October 2009 and February 2010. The statistical analyses were carried out by SPSS version 18.0 and the data were analyzed.

Results: Out of 50 coaches, 94% were males and 6% were females with mean age of 38.74 years. 70% of the coaches found the frequency of injury in the range of 1-5, which included soft tissue injury (46%) followed by bruise and dental trauma. 66% of the coaches believed that boxing was the sports event most likely to cause injury, but boxing, football and rugby (38.5%) were the sports which actually caused injury. The mechanism of injury was mainly due to collision (42%). The protective devices most commonly used were helmet (61.4%) and mouth guards (47.7%). 70% of the coaches felt that non-use of protective devices can cause injuries sometimes. 68% of the coaches found that protective devices enhance the efficiency of the players and 76% considered the use of protective devices based on the type of games.

Conclusions: The results show that there is an increasing acceptance to the use of protective devices among the coaches of Chennai, but its usage was mostly based on the type of game.

Keywords: Coaches; Sports injuries; Protective devices

Introduction

A physically active lifestyle is important for all age groups. Reason to participate in sports and physical activity are many such as pleasure, relaxation, competition, socialization, maintenance and improvement of fitness and health [1]. Participation in sports offers potential benefits for individuals of all ages, such as combating obesity and enhancing cardiovascular fitness. A well-designed exercise program enhances the immediate physical, psychomotor, and intellectual attainments of a child [2]. Thousands of school children play at least one organized sport [3]. It is of growing concern about the injuries that occur during participation in organized sports [4]. Injuries during sport appear to be unavoidable, and up to 30% to 40% of all accidents in

children and adolescents occur during sports [5]. The prevalence of sports injuries vary widely. Comparing injury statistics can be difficult as there is no uniform system for reporting the number of injuries [6,7].

Contact sports are defined as those sports in which players physically interact with each other, trying to prevent the opposing team or person from winning [8]. There is seldom very high incidence of dental trauma compared with all accidents in contact sports varying from 2% to 33% [9,10,11]. Dental and soft tissue injuries are commonly associated with collision and contact sports such as football, ice hockey, soccer, baseball, bicycling, inline skating, gymnastics, basket ball, alpine skiing, softball, volleyball, karate, mountain biking and skateboarding [6,12,13,14,15,16].

Incidence of sports related oral injury vary widely based on the sport played, level of competition, the participant's age and sex [6,17]. Dental injuries are the most common type of orofacial injuries sustained during sports activities [18]. Dental trauma in sports differ from other dental trauma, as it is possible to easily prevent it and there is also a possibility to dramatically reduce the occurrence levels by the use of mouthguards that protect all dental and periodontal structures [19,20]. Dental professionals have long encouraged mouthguard use and dental consultation in organized sports [21]. The only professional sports that require a mouthguard is boxing [22]. In New Zealand, despite a high incidence of avulsive injuries, sports coaches generally had an inadequate knowledge for managing such conditions [23]. Injuries can counter the beneficial effects of sports participation at a young age if a child or adolescent is unable to participate because of the residual effects of injury [24,25].

Some sports medicine professionals pointed directly to poor coaching as a factor in injury [26]. Many young athletes are never taught the proper conditioning methods or technique for their sports [27], as the coaches simply do not have the proper knowledge base for instructing in correct sports techniques [28]. However Smith and Smoll and Smith et al proved the effectiveness of certain communication techniques used by coaches to interact with young athletes during their coaching [29,30,31]. The risks

Materials and Methods

This study is a descriptive survey research design which was carried out between October 2009 and February 2010. The questionnaire which was used in the previous studies on coaches by Lehl G [32] was modified and finally 15 open and closed ended questions were structured. [Table 1] This questionnaire was distributed to fifty coaches who assisted with the study after informed consent. The survey was conducted on the various sports teams in different places of Chennai like YMCA, Radha Krishna stadium, Nehru Park and sports authority of India. The investigators went through this questionnaire with each of the participants separately in order to avoid blanket answers by the entire team of coaches. The questionnaire sought information

1.	Personal details Name: Age: Qualification:
2.	Enumerate the games you coach and please mention the experience you have in the same? <div style="display: flex; justify-content: space-between;">GamesExperience in years</div>
3.	Are sports injuries and their prevention a part of your syllabus? Yes No Not sure
4.	Have you interacted with medical or dental experts regarding safety measures in sports? Yes No
5.	List (in order of priority) the three sports events most likely to cause oro-facial or dental injury? (a) (b) (c)
6.	How often have players under your coaching have sustained an oro-facial or dental injury (in last year)? (a)None (b)1-5 (c)6 or more
7.	What sports event the injury were related to? (a) (b) (c)
8.	What were the mechanisms of injuries? (a) Falls (b) Collision (c) Hit by a ball/hockey stick/hard object (d) Fights between players
9.	What were the types of injuries please tick? (a) Bruise (b) Cut-lip, cheek, tongue (c) Broken tooth, Tooth lost (d) Fracture of facial bones (e) Others
10.	What were the first-aid measures taken for the players? (a) First aid done by me (b) First aid done by any associated medical officer (c) Referral to the hospital/dental office
11.	List the oral or facial devices used in sports that you know to avoid the injuries? <div style="display: flex; justify-content: space-between;">GamesProtective devices</div>

12.	How often were the injuries related to the non use of protective devices? (a) Always (b) Sometimes (c) Never
13.	Do you feel the use of protective devices will have any impact on the efficiency of the players? (a) Reduce efficiency? (b) Enhance efficiency? (c) Any other comment?
14.	In your opinion the usage of protective devices in sports is (a) Always required irrespective of the type of game (b) Required based on the type of game (c) Not required at all
15.	Would you like to recommend any other new form of protective devices that is needed in your sport/game other than mentioned below? (a) Mouthguard, helmet, facemask, knee cap (b)-----

about what sport(s) they coached, the experience they had as a coach, the rate of oral trauma they had seen during sport activities, the mechanism of injuries, type of injury, various protective devices they knew for each games and its usage in sports to avoid injuries.

Statistical Analysis

Descriptive statistics such as mean age and SD were carried out. Graphic presentation of data using the pie chart was also employed. The statistical analyses were carried out using Statistical Package for Social Sciences (SPSS for windows) version 18.0 and the data were analyzed as percentages and proportions.

Results

A total of 50 coaches completed the questionnaire of which 94% were males and 6% were females. The mean age of the coaches was 38.74 years. 50% of the coaches had a Physical Education Training (PET) and 34% of the coaches had 11 or more years of coaching experience. The games for which coaching was provided is listed in (Table 2), which showed that football was most common (20%). 96% of the coaches had sports injury prevention as part of their syllabus and 80% of the coaches had interacted with the medical or dental experts regarding the safety measures in sports.

70% of the coaches found the frequency of injury in the range of 1-5 [Table 3]. 66% of the coaches believed that boxing was the sports event most likely to cause oro-facial injury (Table 1). Boxing again stood first among the sports events (38.5%) along with football and rugby which actually caused injury (Table 2). In majority of the injuries, the mechanism by which they were caused was due to collision (42%) (Table 3). The most common injury was a soft tissue injury (46%) like cut lip, cheek or tongue, followed by bruise and dental trauma (Table 4). 46.20% of the coaches gave first aid by themselves.

On comparing the games and protective devices used, 31.4% of the coaches opted mouthguard as the most preferred protective device for boxing. Helmet was the most preferred protective device for cricket (49%) and hockey (40%). In games like football, rugby and others, the most preferred protective

Table 2: Games for which coaching is provided.

Games for which now coaching is provided	Frequency	Percentage
Athletics	4	8%
Basket ball	5	10%
Boxing	3	6%
Cricket	8	16%
Foot ball	10	20%
Hockey	5	10%
Kabbadi	3	6%
Kho-kho	3	6%
Rugby	4	8%
Volleyball	5	10%
Total	50	100%

Table 3: Frequency of sustaining an oro-facial or dental injury under respondent's coaching.

Frequency of sustaining an injury	Frequency	Percentage
None	11	22%
1-5	35	70%
6 or more	4	8%
Total	50	100%

Table 4: Frequency of oro-facial injury when not using protective devices.

Frequency of injury	Frequency	Percentage
Always	14	28%
Sometimes	35	70%
Never	1	2%
Total	50	100%

devices opted was from the other's category which includes shinguard, abdomen guard etc. (Table 5)

The frequency of injuries when not using protective devices is shown in (Table 4). Of which 70% of the coaches felt that non-use of protective devices can cause injuries sometimes and 28% of the coaches felt that non-use of protective devices can

Table 5: Other Protective devices other than Mouth guards (Multiple answers).

Other Protective devices	Responses	
	N	Percent
Shinguard	9	37.5
Head Gear	2	8.3
Abdomen guards	1	4.2
Armguards	2	8.3
Elbow cap	3	12.5
Kneecap	1	4.2
Crap bandage	3	12.5
Boot / spikes	3	12.5
Total	24	100.0

*(Out of 50 persons only 24 have answered for this question)

cause injuries always. 68% of the coaches found that protective devices enhance the efficiency of the players and 76% of the coaches considered the use of protective devices based on the type of games. Out of 50 coaches only 24 coaches responded for protective devices other than mouthguard, helmet, facemask and knee cap (Table 5).

- Coaches have witnessed 1-5 numbers of oro-facial or dental injuries during their coaching.
- Sports that caused orofacial injuries were boxing, football and rugby.
- Mechanism of injury was collision by players.
- Type of injury witnessed was soft tissue injuries.
- Games - Protective devices used
- Boxing - Mouthguards
- Cricket and hockey - Helmet
- Most of the coaches felt non use of protective devices to cause injury was sometimes not always.
- Even though most of the coaches agreed that protective devices enhance the efficiency of the players, it was used or they advocated its usage based on the game they played.

Discussion

Injury has always been recognized as a natural risk of participation in organized sport. However, many injuries may be preventable, particularly those resulting from a lack of proper, well-supervised training and participation. The present study focused on the knowledge and attitudes of coaches regarding oro-facial injuries and their prevention since they are the main influence on athletes. 70% of the coaches in the present study found the frequency of oro-facial injuries between 1-5 and 70% of these injuries were mostly due to non-use of protective devices. In a similar study by Lehl G, 40% of the coaches found

the frequency of injury between 1-5 [32]. Authors who found the frequency of injury due to non-use of protective devices were 82% by Lehl G, 63% by Garon MW, et al. 75% by Mc Nutt, et al. and 48% by Rob Berg, et al. [32,33,34,35].

Though 66% of the coaches in the present study believed that boxing most likely caused injury, the sports events that actually caused injury along with boxing (39%) were also football (39%) and rugby (39%). This may be because probably higher percentage of coaches were having football as their main sport (20%). Similar results were also found by Mc Nutt, et al. and Garon MW, et al. who found the highest oro-facial injuries with football [33,34]. According to Fos, et al. the high velocity and intensity of football, hockey and rugby place them in a high risk category for oro-facial injuries whereas Kujala, et al. observed that injury rates were higher in those sports involving more frequent and powerful body contact like ice hockey and karate [36,37].

In the present study, majority of the injuries occurred due to collision (42%) followed by hit by an object (32%). The most common type of injury were soft tissue injuries (46%) followed by bruising (24%) and dental trauma (16%). In a similar study, Blinkhorn FA found that falls and collision were the main cause of injury (38). Rob Berg, et al. found in his study that the most common injury was a cut lip, tongue or cheek (82.4%). He also suggested that this type of injury should be a cause for great concern, as they may represent a tooth-object impact in which significant tooth damage was avoided only fortuitously [35].

The protective devices most often preferred by the coaches in the present study were helmet and mouthguards similar to the results by Lehl G [32]. Ranalli DN observed that appropriate and properly fitted protective mouthguards, headgear and helmets prevent oro-facial trauma [39]. Helmets and facemasks when used properly enhance player safety and reduce morbidity [22,40]. It is a promising finding that 80% of the coaches had interacted with medical or dental experts regarding injury prevention and 68% of the coaches found that protective devices enhance the efficiency of the players.

Conclusion

This study identified several important perceptions and knowledge of the coaches in Chennai, India, regarding sports related injury and the usage of various protective devices. Majority of the coaches indicated that they were aware that their trainees were at risk of experiencing oro-facial injuries, including soft tissue injury and dental trauma. It appears that there is an increasing acceptance to the use of protective devices among coaches but its usage was mostly based on the type of game. We feel that preventable injuries do continue to occur since not all players comply with rules, and not all coaches insist that players wear protective devices during practice and in competition. We recommend that the dentist not only involve themselves in giving adequate information to coaches about protective devices, but also counsel patients, especially children, adolescents and young adults, regarding participation in sports so that adequate oral protection can be recommended.

References

1. Bahr R, Holme I. Risk factors for sports injuries: a methodological approach. *Br J Sports Med.* 2003;37:384-392.
2. Shephard RJ. Physical activity and the child health. *Sports Med.* 1984;1(3):205-233.
3. Nowjack-Raymer RE, Gift HC. Use of mouthguards and headgear in organized sports by school-aged children. *Public Health Rep* 1996;111(1):82-86.
4. Micheli Lyle J, Mark DJ. *Sportwise: an essential guide for young athletes, parents, and coaches.* Boston MA: Houghton Mifflin Company;1990.
5. Smith AD, Tao SS. Knee injuries in young athletes. *Clin Sports Med.* 1995;14(3):629-650.
6. Kumamoto DP, Maeda Y. A literature review of sports-related orofacial trauma. *Gen Dent.* 2004;52(3):270-281.
7. Tesini DA, Soporowski NJ. Epidemiology of orofacial sports-related injuries. *Dent Clin North Am.* 2000;44(1):1-18.
8. Dorney B. Dental screening for rugby players in New South Wales, Australia. *FDI World* 1998;7:10-13.
9. Roaas A, Nilsson S. Major injuries in Norwegian football. *Br J Sports Med.* 1979;13(1):3-5.
10. Pritchett JW. Cost of high school soccer injuries. *Am J Sports Med.* 1981;9(1):64-66.
11. National Health and Medical Research Council. *Football injuries of the head and neck.* The Australian Government Publishing Services, 1994:1-128.
12. Soporowski NJ, Tesini DA, Weiss AI. Survey of orofacial sports-related injuries. *J Mass Dent Soc.* 1994;43(4):16-20.
13. Bayliss T, Bedi R. Oral maxillofacial and general injuries in gymnasts. *Injury.* 1996;27(5):353-354.
14. Maestrello-de Moya M, Primrosch R. Orofacial trauma and mouth protector wear among high school varsity basket-ball players. *J Dent Child.* 1989;56(1):36-39.
15. Gassner R, Vazquez Garcia J, Leja W, Stainer M. Traumatic dental injuries and Alpine skiing. *Endod Dent Traumatol.* 2000;16(3):122-127.
16. Lang B, Pohl Y, Filippi A. Knowledge and prevention of dental trauma in team handball in Switzerland and Germany. *Dent Traumatol.* 2002;18(6):329-334.
17. Bijur PE, Trumble A, Harel Y, Overpeck MD, Jones D, Scheidt PC. Sports and recreation injuries in US children and adolescents. *Arch Pediatr Adolesc Med.* 1995;149(9):1009-1016.
18. Scott J, Burke FJT, Watts DC. A review of dental injuries and the use of mouthguards in contact team sports. *Br Dent J.* 1994;176(8):310-314.
19. Andreasen JO, Andreasen FM. *Textbook and color atlas of traumatic injuries to the teeth*, 3rd ed. Munksgaard, Copenhagen.1994;719-735.
20. Kerr IL. Mouthguards for the prevention of injuries in contact sports. *Sports Med.* 1986;3(6):415-427.
21. Powers JM, Godwin WC, Heintz WD. Mouth protectors and sports team dentists. *JADA.* 1984;109(1):84-7.
22. Ranalli DN. Prevention of craniofacial injuries in football. *Dent Clin North Am.* 1991;35(4):627-645.
23. Stokes AN, Anderson HK, Cowan TM. Lay and professional knowledge of methods for emergency management of avulsed teeth. *Endod Dent Traumatol.* 1992;8(4):160-162.
24. Caine DJ, Maffulli N. Epidemiology of children's individual sports injuries. An important area of medicine and sport science research. *Med Sports Sci.* 2005;48:1-7.
25. Erlandson MC, Sherar LB, Mirwald RL, Maffulli N, Baxter-Jones AD. Growth and maturation of adolescent female gymnasts, swimmers, and tennis players. *Med Sci Sports Exerc.* 2008;40(1):34-42.
26. Murphy P. Youth sports coaches: using hunches to fill a blank page. *Physician Sportsmed* 1985;13(4):136-142.
27. Micheli LJ. Sports injuries in the young athlete: questions and controversies In: Micheli LJ, ed. *Pediatric and Adolescent Sports Medicine.* Boston. MA: Little Brown and Co: 1984:1-8.
28. Quain RJ. An overview of youth coaching certification programs. *Adolescence.* 1989;24:541-547.
29. Smith RE, Smoll FL. Self-esteem and children's reactions to youth sports coaching behaviors: A field study of self-enhancement processes. *Dev Psychol.* 1990;26:987-993.
30. Smith RE, Smoll FL, Curtis B. Coaching behaviors in Little League baseball. In: Smoll FL, Smith RE, eds. *Psychological Perspectives in Youth Sports.* Washington, DC: Hemisphere; 1978:173-201.
31. Smith RE, Smoll FL, Curtis B. Coach effectiveness training a cognitive behavioral approach to enhancing relationship skills in youth sport coaches. *J Sport Psychol.* 1979;1:59-75.
32. Lehl G. Perceptions of Chandigarh sports coaches regarding orofacial injuries and their prevention. *J Indian Soc Pedod Prev Dent.* 2005;23(2):67-70.
33. Garon MW, Merkle A, Wright JT. Mouth protectors and oral trauma: a study of adolescent football players. *JADA.* 1986;112:663-665.
34. McNutt T, Shannon SW Jr, Wright JT, Feinstein RA. Oral trauma in adolescent athletes: a study of mouth protectors. *Pediatr Dent.* 1989;11(3):209-213.
35. Rob Berg, Berkey DB, Tang JMW, Altman DS, Londeree KA. Knowledge and attitudes of Arizona high-school coaches regarding oral-facial injuries and mouthguard use among athletes. *JADA.* 1998;129:1425-1432.
36. Fos PJ, Pinkham JR, Ranalli DN. Prediction of sports related dental traumatic injuries. *Dent Clin North Am.* 2000;44:19-33.
37. Kujala UM, Taimela S, Antti-Poika I, Orava S, Tuomian R, Myllynen P. Acute injuries in soccer, ice hockey, volleyball, basketball, judo and karate. Analysis of national registry data. *BMJ.* 1995;311:1465-1468.
38. Blinkhorn FA. The aetiology of dento-alveolar injuries and factors influencing attendance for emergency care of adolescents in the North West of England. *Endod Dent Traumatol.* 2000;16:162-165.
39. Ranalli DN. Prevention of sports-related traumatic dental injuries. *Dent Clin North Am.* 2000;44:35-51.
40. Rontal E, Rontal M. Maxillofacial injuries in football players, an evaluation of current facial protection. *J Sports Med Phys Fitness.* 1971;11(4):241-245.