

Alternative Source of Protein “Silkworm Pupae” (*Bombyx Mori*) In Coldwater Aquaculture

Oyas A Asimi^{1*}, Tariq H Bhat², Nasir H³ and Irfan K⁴

¹Assistant Professor, Fish Nutrition, Division of Fish Nutrition and Biochemistry, Faculty of Fisheries, Rangil-190 006, SKUAST-K, India

²Assistant Professor, Division of Post Harvest Technology, Faculty of Fisheries, Rangil-190 006, SKUAST-K, India

³Assistant Professor, Division of Social Science, Faculty of Fisheries, Rangil-190 006, SKUAST-K, India

⁴Assistant Professor, Division of Fish Breeding and Genetics, Faculty of Fisheries, Rangil-190 006, SKUAST-K, India

Received: November 08, 2017; Accepted: November 20, 2017; Published: November 29, 2017

*Corresponding author: Oyas A Asimi, Assistant Professor, Fish Nutrition, Division of Fish Nutrition and Biochemistry, Faculty of Fisheries, Rangil-190 006, SKUAST-K, India, E-mail: asimioa@gmail.com

Abstract

The main constraint presently is the availability of sufficient/required quantity of cost effective quality feed. At present one of the main fish feed ingredient (fishmeal) is procured from outside the state due to which the cost of production is very high. In order to make available sufficient quantity of cost effective feed and cater to the demand of various rearing units for feed in the State, it is high time to replace fishmeal with alternate source of protein. Kashmir valley produces good quantities of feed materials derived from crops and other sources. These include a wide variety of oil-seed cakes and meals, pulses, and mill by-products of seeds and grains. Also available are appreciable quantities of by-products from the meat, fish, fruit processing industries and particularly silkworm pupae. Keeping in view the nutritional value of silkworm pupae and its profuse availability in the state, cost effective feed could be formulated and feeding trials in coldwater aquaculture could be conducted. Lot of research has been done on the replacement of fishmeal throughout the world.

Key words: Fishmeal; Silkworm; Aquaculture;

Introduction

Silkworms have a crunchy exterior and reportedly, the taste and texture of mashed potato inside when fresh! A silkworm pupa is the by-product after the silk thread has been wound off from the cocoon and can serve as feedstuff. The most common species for commercial silk production is the mulberry silk worm (*Bombyx mori*) of the sub-phylum Tracheata (articulated animals). The caterpillar of the moth feed chiefly on the mulberry leaf. The percentages of total protein and lipid contents by dry weight are 55.6 and 32.2% respectively. Silkworm pupae protein had high levels of essential amino acids such as valine, methionine and phenylalanine. The contents of essential amino acids in silkworm pupae protein satisfied the FAO/WHO/UNU suggested requirements (2007).

Sericulture in Kashmir

The history of sericulture in J&K State is traced, and the silk industry was demonopolised in 1988. About 27,000 farming families produce around 829 MT of cocoons worth Rs.5.50 crore. An increase to around 1,500 MT is planned in the next decade. Productivity and quality need to be improved through better packages of mulberry cultivation and silkworm rearing, and high yielding quality silkworm breeds. Majority of cocoon production coming from the spring crop - an autumn crop has been introduced but yield and quality are not yet good.

Major input cost in aquaculture is the running cost expended on feeding the cultured fish.

Protein is the most expensive component in fish feeds. Fish meal is most important feed ingredient as dietary protein source for cultured fish. Shortage and high cost of fish meal make fish feed expensive hence the search for alternative sources of protein in fish feeds. Generally, the feed stuffs of animal origins are considered better alternative protein source in et.al fish diets because of their higher content and other superior indispensable amino acids than the plant origins [23]. Several animal protein sources were evaluated to formulate the diets for fish such as poultry by-product meal, meat and bone meal, snail and other invertebrate meal [10, 13, 21, 25, 33]. Main tendency is to partial replacement of fish meal with alternative protein source [30, 31]. Insects may constitute a significant biomass (Table 1).

Nutritional attributes

Silkworm (*Bombyx mori*) pupa is a protein-rich feed ingredient with a high nutritional value. Its crude protein content ranges from 50% DM to more than 80% DM (for defatted meal). The lysine (6-7% of the protein) and methionine (2-3% of the protein) contents are particularly high. However, the true protein

Table 1: Cocoon production during last few years (Source: Sericulture Development Department, Jammu and Kashmir, 2016)

Year	Cocoon Production (MT)	Income generation (Lac Rs.)	Avg. Price for "A" grade (Rs.)	Productivity per Oz. (Kg)	Rearers population (Nos.)
2008-09	738	455.67	192.00	31.00	19700
2009-10	810	800.00	300.00	32.00	22800
2010-11	860	1100.00	410.00	32.5	25500
2011-12	917	963.00	350.00	33.00	27000
2012-13	901	1193.00	397.00	32.00	28000
2013-14	1022	2226.00	650.00	34.23	29400
2014-15 (Spring)	1105 (P)	-	750.00 (P)	42.00 (P)	31882

(calculated as the sum of amino acids) in silkworms was found to correspond to only 73% of the crude protein content (Finke, 2002), which was explained by the presence of chitin, since this component contains nitrogen. However, the chitin content of pupae meal is relatively low, about 3-4% DM [6, 28]. The presence of chitin and insoluble protein may also explain the presence of fibre, and values of 6-12% DM of ADF have been reported [6, 11]. Un-defatted pupae meal is rich in fat, typically in the 20-40% DM range. Defatted meal contains less than 10% oil in the DM. Silkworm oil contains a high percentage of polyunsaturated fatty acids, notably linolenic acid (18:3), with reported values ranging from 11 to 45% of the total fatty acids [11,20,32]. Compared to other animal by-products, silkworm pupae meal is relatively poor in minerals [3-10% DM]. Silkworm litter appears to have an extremely variable composition, with crude protein values reported to be between 15 and 58% DM [3, 32].

The benefits of silkworm in fish feed

Silkworm pupae in a fish diet are a great way to simulate the type of nutrient dense variety that Koi and other aquarium fishes would experience if they were kept in mud pond. A fish in the wild graze on a wide variety of edible plants, algae and insects throughout the day. In order to replicate that varied food source in a tank or pond, including some other nutrient sources, like silkworm pupae, is a great idea.

While there are a number of different silkworm pupae options available on the market, Hikari® manufactures a unique product which is processed and packaged to avoid oxidation, a common issue with competitive silkworm pupae type products. This fresher and more potent delivery system of for these nutrients help fish avoid potential health problems from too much of a good thing or rancidity.

Use of silkworm pupae meal in aquaculture

When silkworm meal replaced part, or all, of the fish meal in common carp (*Cyprinus carpio*) diets, similar performance (growth and feed conversion) was observed [22]. Un-defatted silkworm pupae meal was safely used up to 50% without hampering growth and meat quality of the fish (Nandeesh et al., 2000). In a comparison between silkworm pupae meal and plant leaf meals (alfalfa and mulberry), feed conversion efficiency,

nutrient digestibility and nutrient retention were better for diets based on silkworm meal than for diets based on plant leaf meals [29].

In silver barb fingerlings (*Barbonymus gonionotus*), highest growth rates were observed in fish fed a diet with about 38% of total dietary protein replaced by silkworm pupae meal [15].

Mahseer fingerlings (*Tor khudree*) fed a diet containing 50% defatted silkworm pupae at 5% body weight had better growth and survival rates than fingerlings fed no or lower amounts of silkworm pupae [24].

Fermented silkworm pupae silage or untreated fresh silkworm pupae paste were incorporated in carp feed formulations, replacing fish meal, in a polyculture system containing the Indian carp (*Catla catla*), mrigal carp (*Cirrhinus mrigala*), rohu (*Labeo rohita*) and silver carp (*Hypophthalmichthys molitrix*). Survival rate, feed conversion ratio and specific growth rate were better for fermented silkworm pupae silage than for untreated silkworm pupae or fish meal (Rangacharyulu et al., 2003). In rohu, undefatted silkworm pupae and defatted silkworm pupae produced significantly better protein digestibility values than fish meal [8].

Mozambique tilapia (*Oreochromis mossambicus*) were able to utilize the protein of both defatted and undefatted silkworm meal (apparent protein digestibility of 85-86%) [8].

Chum salmon fry (*Onchorhynchus keta*) fed over 6 week diets supplemented with 5% silkworm pupae meal at the expense of fish meal did not show improvement in growth rate and protein content, though silkworm supplementation enhanced feed efficiency [1].

Silkworm pupae meal was used as a substitute for fish meal at up to 75% of the protein in Asian stinging catfish (*Heteropneustes fossilis*) diets without adverse effects on growth [8].

Un-defatted silkworm pupae were found to be a suitable fish meal substitute in diets for walking catfish (*Clarias batrachus*). Digestibility of the crude protein in silkworm meal was found to be similar to that of fish meal [2].

In Japanese sea bass (*Lateolabrax japonicus*), the energy digestibility of un-defatted silkworm pupae meal (73%) was

lower than that of poultry by-product meal, feather meal, blood meal and soybean meal, but comparable to that of meat and bone meal. Protein digestibility was also lower (85%) than that of poultry by-product meal, blood meal and soybean meal but it was comparable to that of feather meal and higher than that of meat and bone meal [12].

Shrimp growth trials showed that digestive efficiency was reduced when silkworm meal was used to replace fish meal [26].

In juvenile abalones (*Haliotis discus hannai* Ino.), a combination of soybean meal and silkworm pupae meal replaced fish meal, resulting in slightly higher survival rate and better growth performance (Cho SungHwoan, 2010).

Antioxidant activity of silkworm pupae oil

Supanida Winitchai et al. extracted oil from five native Thai silkworm varieties. The oil extracted by the Soxhlet method from silkworm varieties showed free radical scavenging activity [27]. Deori et al. investigated antioxidant activity of pupae of the muga and Eri silkworm and concluded that, the pupae could be used as natural antioxidants on food products [5].

Conclusion

Silkworm pupae protein had high levels of essential amino acids such as valine, methionine and phenylalanine. The contents of essential amino acids in silkworm pupae protein satisfied the FAO/WHO/UNU suggested requirements (2007). In addition, they also possessed n-3 fatty acids, especially α -linolenic acid (36.3%), as a major component. Keeping in view the above facts, silkworm pupae can be used as an alternative source of protein in fish feed instead of fishmeal.

References

- Akiyama T, Murai T, Hirasawa Y, Nose T. Supplementation of various meals to fish diet for chum salmon fry. *Aquaculture*. 1984;37(3):217-222. Doi.org/10.1016/0044-8486(84)90154-6
- Borthakur S, K. Sarma. Protein and fat digestibility of some non-conventional fish meal replacers incorporated in the diets of fish *Clarias batrachus* (Linn.). *Environ Ecol*. 1998;16(2):368-371. Doi: 10.1016/S0550-3213(00)00030-4
- Chen Yao wang. Animal raising and plant cultivation on an integrated fish farm. *Integrated fish farming in China*. 2003.
- Cho Sung Hwoan. Effect of fishmeal substitution with various animal and/or plant protein sources in the diet of the abalone *Haliotis discus hannai* Ino. *Aquacult Res*. 2010;41(10):e587-e593. DOI: 10.1111/j.1365-2109.2010.02561.x
- Deori M, Dipali Devi, Rajalakshmi Devi. Nutrient composition and antioxidant activities of muga and eri silkworm pupae. *International Journal of Science and Nature*. 2014;5(4):636-640.
- Finke M D. Complete nutrient composition of commercially raised invertebrates used as food for insectivores. *Zoo Biology*. 2002;21(3):269-285. DOI: 10.1002/zoo.10031
- Hikari. Making Species Specific Diets For Decades! "Check out Silkworm Selects™ – a great way to introduce this ancient nutrient booster to your tank or pond pets.
- Hossain M A, N Nahar, M Kamal, M N Islam. Nutrient digestibility coefficients of some plant and animal proteins for tilapia (*Oreochromis mossambicus*). *J Aquacult Trop*. 1992;7(2):257-265.
- Hossain M A, N Nahar, M. Kamal. Nutrient digestibility coefficients of some plant and animal proteins for rohu (*Labeo rohita*). *Aquaculture*. 1997;151(1-4):37-45. Doi.org/10.1016/S0044-8486(96)01481-0
- Hu M, Wang Y, Wang Q, Zhao M, Xiong B, Yujiang Zhao, et al. Replacement of fish meal by rendered animal protein ingredients with lysine and methionine supplementation to practical diets for gibel carp, *Carassius auratus gibelio*. *Aquaculture*. 2008;275:260-265. DOI: 10.1016/j.aquaculture.2008.01.005
- Ioselevich M, Steingaß H, Rajamurodov Z, Drochner W. Nutritive value of silkworm pupae for ruminants. *VDLUFA Kongress. Qualitätssicherung in landwirtschaftlichen Produktionssystemen*. Rostock. 2004;116-108.
- Ji WenXiu, Wang Yan and Tang JinYu. Apparent digestibility coefficients of selected feed ingredients for Japanese sea bass (*Lateolabrax japonicus*) reared in sea water. *J Fish China*. 2010;34(1):101-107.
- Li K, Wang Y, Zheng ZX, Jiang RL, Xie NX, Dominique P Bureau. Replacing fish meal with rendered animal protein ingredients in diets for Malabar grouper, *Epinephelus malabaricus*, reared in net cages. *J World Aquacult Soc*. 2009;40(1):67-75. DOI: 10.1111/j.1749-7345.2008.00227.x
- MA Khan, MN Ahmad, SB Zeya. *Sericulture in Jammu and Kashmir - Past, present and future*. 2003.
- Mahata SC, Bhuiyan AMA, Zaheer M, Hossain MA, Hasan MR. Evaluation of silkworm pupae as dietary protein source for Thai Sharpunti, *Puntius gonionotus* (Bleaker). *Journal of Aquaculture Trop*. 1994;9(1):77-85.
- Muthuraju, Shruthi PA. Review – Bionutritional Science of Silkworm Pupal residue to Mine New ways for utilization. *Int J Adv Res Biol Sci*. 2015;2(9):135-140.
- Nandeesh MC, Gangadhara B, Varghese TJ, Keshavanath P. Growth response and flesh quality of common carp, *Cyprinus carpio* fed with high levels of non-defatted silkworm pupae. *Asian Fish Sci*. 2000;13:235-242.
- Patil RR, Bhattacharyya A, Rayar SG, Uppar V, 2013. Gold-It's Effect on Cocoon and Silk traits of Mulberry silkworm, *Bombyx mori* L, 6th BACSA International conference "Building Value Chains in Sericulture" "BISERICA" 2013.
- Rangacharyulu PV, SS Giri, BN Paul, KP Yashoda, RJ Rao, NS Mahendrakar, SN Mohanty, et al. Utilization of fermented silkworm pupae silage in feed for carps. *Bioresour Technol*. 2003;86(1):29-32.
- Rao UP. Chemical composition and nutritional evaluation of spent silkworm pupae. *Am Chem Soc*. 1994;42(10):2201-2203.
- Rawles SD, Thompson KR, Brady YJ, Metts LS, AksoyMY. Effects of replacing fish meal with poultry byproduct meal and soybean meal and reduced protein level on the performance and immune status of pond-grown sunshine bass (*Morone chrysops* × *M. saxatilis*). *Aquacult Nutr*. 2011;17(3):708-721. DOI: 10.1111/j.1365-2095.2010.00831.x
- Rehman, MA, M Zaher, MA Mazed, MZ Haque and SC Mahata. Replacement of costly fishmeal by silkworm pupae in diet of mirror carp. *Pak J Sci and Industrial Res*. 1997.39(1-4):64-67.
- Robinson EH, Li MH. Comparison of practical diets with and without animal protein at various concentrations of dietary protein on performance of channel catfish *Ictalurus punctatus* raised in earthen

- ponds. J World Aquaculture Society. 1998;29(3):273-280. DOI: 10.1111/j.1749-7345.1998.tb00647.x
24. Shyam Sunder, M Mohan, HS Raina, R Singh, RS Haldar. Culture of golden mahseer, *Tor putitora* (Ham.) in Kumaon Himalaya. 1. Mass scale production of stocking material. In: Proceedings of Third Indian Fisheries Forum. 1993.
25. Sogbesan AO, Ugwumba AAA, Madu CT. Nutritive potentials and utilization of garden snail (*Limicolaria aurora*) meat meal in the diet of *Clarias gariepinus* fingerlings. Afr J Biotechnol. 2006;5(20):1999-2003.
26. Sumitra-Vijayaraghavan, Wafar MVM, Royan JP. Ind J. 1978;7:195-197.
27. Supanida Winitchai, Jiradej Manosroi, Masahiko Abe, Korawinwich Boonpisuttinant, Aranya Manosroi. Free Radical Scavenging Activity, Tyrosinase Inhibition Activity and Fatty Acids Composition of Oils from Pupae of Native Thai Silkworm (*Bombyx mori* L.). J Nat Sci. 2011;45:404-412.
28. Suresh H N, Mahalingam C A, Pallavi. Amount of chitin, chitosan and chitosan based on chitin weight in pure races of multivoltine and bivoltine silkworm pupae *Bombyx mori* L. Int J Sci Nature. 2012;3(1):214-216.
29. Swamy HVV, KV Devaraj. Nutrient utilization by common carp (*Cyprinus carpio* Linn) fed protein from leaf meal and silkworm pupae meal based diets. Indian J Anim Nutr. 1994;11(2):67-71.
30. Tacon AG. Feed ingredients for warm water fish: fishmeal and other feedstuffs. FAO Circular no 856, FAO Rome, 64.
31. Tacon AG, Akiyama DM. Feed ingredients. In: Crustacean nutrition (ed. D'Abramo LR., Conklin DE, Akiyama, DM), Advances in world aquaculture, World Aquaculture Society, Baton Rouge. 6:411-472.
32. Usub T, Lertsatitthanakorn C, Poomsaad, L Wiset, L Yang, Siriamornpun S. Experimental performance of a solar tunnel dryer for drying silkworm pupae. Biosyst Eng. 2008;101(2):209-216. Doi. org/10.1016/j.biosystemseng.2008.06.011
33. Xue M, Zhou Z, Wu XF, Yu Y, Ren ZL. Partial or total replacement of fish meal by meat and bone meal in practical diets for Nile tilapia, *Oreochromis niloticus*. ACTA Zoonutri Sinica. 2003;1:106-112.